

MONOTOPE [™] product information for

Respiratory Syncytial Virus

I. Monoclonal Antibody (Mouse) Specificity

Product No.'s	Ig class	
0611	NA	IgG1 specific for the nucleoprotein of RSV, performs well in indirect IFA & ELISA. Broad reactivity.
0631	NA	IgG2a specific for the fusion protein of RSV, types A&B. Neutralizes. Pairs well in ELISA*. Functions in IFA.
0641	NA	IgG1 specific for the fusion protein of RSV, types A&B. Neutralizes. Pairs well in ELISA*. Functions in IFA.
0651	NA	IgG2a specific for the fusion protein of RSV, types A&B. Neutralizes. Pairs well in ELISA*. Functions in IFA.
0661	0663	IgG1 specific for the nucleoprotein of RSV, performs well in indirect IFA. Pairs with #0604 in ELISA.
0671	0673	IgG2a specific for the nucleoprotein of RSV, performs well in indirect IFA & ELISA
0681	NA	IgG2b specific for the fusion protein of RSV, functions in indirect IFA & ELISA. Neutralizes

II. Purified Preparations

- Product No.'s
- 0611**
 - 0631
 - 0641**
 - 0651**
 - ~~0661~~
 - 0671
 - 0681

MONOTOPE™ purified preparations consist of >90% pure mouse monoclonal antibody which has been purified from ascites fluid or culture medium by protein A chromatography or sequential differential precipitations. The final preparation is formulated to a protein concentration of 100 µg/ml in 0.01 M phosphate buffered saline, pH 7.2 and contains 0.1% sodium azide. Each vial contains 1.0 ml. This product contains no stabilizing proteins and should be stored at -20°C until ready for use. Avoid repeated freeze-thawing by storing multiple aliquots at -20°C. Working dilution must be determined by the user. Suggested starting ranges are 1:10-1:50 for IFA, blotting & IHC and 1:20-1:200 for ELISA. Custom conjugation of this antibody with HRP, alkaline phosphatase, biotin and fluorescein is available on a minimum order basis

THESE PRODUCTS ARE FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES

III. Fluorescein Conjugates

- Product No.'s
- NA
 - NA
 - NA
 - NA
 - ~~0663~~
 - 0673
 - NA

These MONOTOPE™ products consist of purified monoclonal antibody conjugated with high purity isomer I of fluorescein isothiocyanate. Care is taken to ensure complete removal of any free fluorescein from the final product. The final preparation is formulated to an antibody concentration of 100 µg/ml in 0.01 M phosphate buffered saline, pH 7.2 containing 0.1% sodium azide plus bovine serum albumin at 10 mg/ml. Each vial contains 1.0 ml. This product should be stored at -20°C until ready for use. Avoid repeated freeze-thawing by storing multiple aliquots at -20°C. Applications for these products include direct FA staining of target antigen in a permissive tissue culture system. Working strength must be determined by the user for each specific application but a starting range of 1:5 - 1:20 is recommended. Acetone fixation of the antigen source is recommended prior to staining.

THESE PRODUCTS ARE FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES

Comments: *See table on back for additional information including optimum pairing in ELISA.