

MONOTOPE™ product information for

Chlamydia trachomatis

I. Monoclonal Antibody (Mouse) Specificity

Product No.'s	Ig class	
1644 NA	IgG2b	Highest affinity for LPS. ELISA pairing below. Genus specific (C. trachomatis, C. pneumoniae & C. psittaci)
1645 NA	IgG2b	High affinity for LPS. ELISA pairing below. Genus specific (C. trachomatis, C. pneumoniae & C. psittaci)
1646 NA	IgG2b	High affinity for LPS. ELISA pairing below. Genus specific (C. trachomatis, C. pneumoniae & C. psittaci)
1647 NA	IgG2a	High affinity for LPS. ELISA pairing below. Genus specific (C. trachomatis, C. pneumoniae & C. psittaci)
1681 1683	IgG1	LPS, Genus specific*, crossreactive with C. pneumoniae & C. psittaci. Functions in IFA, ELISA & IHC
1685 NA	IgG1	LPS, Genus specific*, crossreactive with C. pneumoniae & C. psittaci. Functions in ELISA, IFA & IHC
1671 1673	IgG2a	Specific to MOMP. Recognizes 15 serovars of C. trachomatis. Functions in ELISA, IFA & IHC

II. Purified Preparations

Product No.'s
1644
1645
1646
1647
1681
1685
1671

MONOTOPE™ purified preparations consist of >90% pure mouse monoclonal antibody which has been purified from ascites fluid or culture medium by protein A chromatography or sequential differential precipitations. The final preparation is formulated to a protein concentration of 100 µg/ml in 0.01 M phosphate buffered saline, pH 7.2 and contains 0.1% sodium azide. Each vial contains 1.0 ml. This product contains no stabilizing proteins and should be stored at -20°C until ready for use. Avoid repeated freeze-thawing by storing multiple aliquots at -20°C. Working dilution must be determined by the user. Suggested starting ranges are 1:10-1:50 for IFA, blotting & IHC and 1:20-1:200 for ELISA. Custom conjugation of this antibody with HRP, alkaline phosphatase, biotin and fluorescein is available on a minimum order basis

THESE PRODUCTS ARE FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES

III. Fluorescein Conjugates

Product No.'s
NA
NA
NA
NA
1683
NA
1673

These MONOTOPE™ products consist of purified monoclonal antibody conjugated with high purity isomer I of fluorescein isothiocyanate. Care is taken to ensure complete removal of any free fluorescein from the final product. The final preparation is formulated to an antibody concentration of 100 µg/ml in 0.01 M phosphate buffered saline, pH 7.2 containing 0.1% sodium azide plus bovine serum albumin at 10 mg/ml. Each vial contains 1.0 ml. This product should be stored at -20°C until ready for use. Avoid repeated freeze-thawing by storing multiple aliquots at -20°C. Applications for these products include direct FA staining of target antigen in a permissive tissue culture system. Working strength must be determined by the user for each specific application but a starting range of 1:5 - 1:20 is recommended. Acetone fixation of the antigen source is recommended prior to staining.

THESE PRODUCTS ARE FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES

Comments: * Reactive with all serovars tested (8).

Best LPS ELISA pairs (Capture/Conjugate): 1645/1644, 1644/1644, 1646/1644, 1647/1644