

MONOTOPE™ product information for
Human Papilloma viruses

I. Monoclonal Antibody (Mouse) Specificity

Product No.'s	Ig class	
5011 NA	IgG1	Specific the capsid (L1) of HPV type 6 . Functions in ELISA. Negative in western blot immunoassay.
5021	IgG1	Specific the capsid (L1) of HPV type 11. Functions in ELISA. Negative in western blot immunoassay.
5031	IgG2a	Specific the capsid (L1) of HPV type 16. Functions in ELISA. Negative in western blot immunoassay.
5041	IgG1	Specific the capsid (L1) of HPV type 18. Functions in ELISA. Negative in western blot immunoassay.
5051 NA	IgG2a	Specific the E7 protein of HPV types 16 & 18 . Functions in ELISA & WB. Pairs with #5061.
5061 NA	IgG2b	Specific the E7 protein of HPV types 16 & 18 . Functions in ELISA & WB. Pairs with #5051.

II. Purified Preparations

Product No.'s	5011
	5021
	5031
	5041
	5051
	5061

MONOTOPE™ purified preparations consist of >90% pure mouse monoclonal antibody which has been purified from ascites fluid or culture medium by protein A chromatography or sequential differential precipitations. The final preparation is formulated to a protein concentration of 100 µg/ml in 0.01 M phosphate buffered saline, pH 7.2 and contains 0.1% sodium azide. Each vial contains 1.0 ml. This product contains no stabilizing proteins and should be stored at 2-8°C until ready for use. Avoid repeated freeze-thawing by storing multiple aliquots at -20°C. Working dilution must be determined by the user. Suggested starting ranges are 1:10-1:50 for IFA, blotting & IHC and 1:20-1:200 for ELISA. Custom conjugation of this antibody with HRP, alkaline phosphatase, biotin and fluorescein is available on a minimum order basis

THESE PRODUCTS ARE FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES

III. Fluorescein Conjugates

Product No.'s	NA
	NA
	NA

These MONOTOPE™ products consist of purified monoclonal antibody conjugated with high purity isomer I of fluorescein isothiocyanate. Care is taken to ensure complete removal of any free fluorescein from the final product. The final preparation is formulated to an antibody concentration of 100 µg/ml in 0.01 M phosphate buffered saline, pH 7.2 containing 0.1% sodium azide plus bovine serum albumin at 10 mg/ml. Each vial contains 1.0 ml. This product should be stored at 2-8°C until ready for use. Avoid repeated freeze-thawing by storing multiple aliquots at -20°C. Applications for these products include direct FA staining of target antigen in a permissive tissue culture system. Working strength must be determined by the user for each specific application but a starting range of 1:5 - 1:20 is recommended. Acetone fixation of the antigen source is recommended prior to staining.

THESE PRODUCTS ARE FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES

Comments: