

MONOTOPE™ product information for

Epstein-Barr Virus

I. Monoclonal Antibody (Mouse) Specificity

Product No.'s	Ig class	
0212	NA	IgG2a specific for EBNA-1 protein. Reactive with native antigen*. Functions in ELISA and WB.
0235	NA	IgG1 specific for the early antigen (restricted) homologue Bcl-2. Functions in ELISA, WB, IHC & IP.
0221	0223	IgG1 specific for envelope glycoprotein complex 250/350. Functions in IFA and blotting.
0231	NA	IgG1 specific for viral capsid antigen (VCA); functions well in IFA but not western blot.
0256	NA	IgG2 specific for the early antigen (BAM HI Z) zeta protein. Functions in ELISA, WB, IHC & IP.
0261	NA	IgG2 specific for the " Early antigen-diffuse" (EA-D)of EBV; functions in IFA, western blot & IHC (weak).

II. Purified Preparations

Product No.'s	0212
	0235
	0221
	0231
	0256
	0261

MONOTOPE™ purified preparations consist of >90% pure mouse monoclonal antibody which has been purified from ascites fluid or culture medium by protein A chromatography or sequential differential precipitations. The final preparation is formulated to a protein concentration of 100 µg/ml in 0.01 M phosphate buffered saline, pH 7.2 and contains 0.1% sodium azide. Each vial contains 1.0 ml. This product contains no stabilizing proteins and should be stored at 2-8°C until ready for use.

Working dilution must be determined by the user. Suggested starting ranges are 1:10-1:50 for IFA and 1:20-1:200 for ELISA.

THESE PRODUCTS ARE FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES

III. Fluorescein Conjugates

Product No.'s	NA
	NA
	0223
	NA
	NA
	NA

These MONOTOPE™ products consist of purified monoclonal antibody conjugated with high purity isomer I of fluorescein isothiocyanate. Care is taken to ensure complete removal of any free fluorescein from the final product. The final preparation is formulated to an antibody concentration of 100 µg/ml in 0.01 M phosphate buffered saline, pH 7.2 containing 0.1% sodium azide plus bovine serum albumin at 10 mg/ml. Each vial contains 1.0 ml. This product should be stored at 2-8°C until ready for use. Avoid repeated freeze-thawing by storing multiple aliquots at -20°C. Applications for these products include direct FA staining of target antigen in a permissive tissue culture system. Working strength must be determined by the user for each specific application but a starting range of 1:5 - 1:20 is recommended. Acetone fixation of the antigen source is recommended prior to staining.

THESE PRODUCTS ARE FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES

Comments: * Epitope lies between aa 408-641