

MabyRab™ product information for
Group A Streptococcus

I. Monoclonal Antibody (Rabbit) Specificity

Product No.'s	Ig class	
0435	CHO IgG1	Specific to the Group A Step carbohydrate. Negative vs. other Strep groups*. Pairs in lateral flow appl. ^
0436	HEK293 IgG1	Specific to the Group A Step carbohydrate. Negative vs. other Strep groups*

II. Purified Preparations

Product No.'s	0435
	0436

RabMAB® purified preparations consist of >90% pure rabbit monoclonal antibody which has been purified from culture medium by protein A chromatography or sequential differential precipitations. The final preparation is formulated to a protein concentration of 100 µg/ml in 0.01 M phosphate buffered saline, pH 7.2 and contains 0.1% sodium azide. Each vial contains 1.0 ml. This product contains no stabilizing proteins and should be stored at 2-8°C until ready for use.

Working dilution must be determined by the user. Suggested starting ranges are 1:10-1:50 for IFA and 1:20-1:200 for ELISA.

THESE PRODUCTS ARE FOR RESEARCH USE ONLY NOT FOR USE IN DIAGNOSTIC PROCEDURES

III. Fluorescein Conjugates

Product No.'s	CHO
	HEK293

These MONOTOPE™ products consist of purified monoclonal antibody conjugated with high purity isomer I of fluorescein isothiocyanate. Care is taken to ensure complete removal of any free fluorescein from the final product. The final preparation is formulated to an antibody concentration of 100 µg/ml in 0.01 M phosphate buffered saline, pH 7.2 containing 0.1% sodium azide plus bovine serum albumin at 10 mg/ml. Each vial contains 1.0 ml. This product should be stored at 2-8°C until ready for use. Avoid repeated freeze-thawing by storing multiple aliquots at -20°C. Applications for these products include direct FA staining of target antigen in a permissive tissue culture system. Working strength must be determined by the user for each specific application but a starting range of 1:5 - 1:20 is recommended. Acetone fixation of the antigen source is recommended prior to staining.

THESE PRODUCTS ARE FOR RESEARCH USE ONLY NOT FOR USE IN DIAGNOSTIC PROCEDURES

Comments: * No cross reactivity with: Streptococci groups B, C, D, F & G + other bacteria.

^ This Rabbit Monoclonal antibody pairs in lateral flow with: itself, rabbit poly 1511 and goat poly 1501.