

Immunochemicals for Infectious Disease Research www.ViroStat-Inc.com

P.O. Box 8522 Portland, Maine 04104 207-856-6620 207-856-6864 (fax)

RabMAB[®]

product information for

Group B Streptococcus

I. Monoclonal Antibody (Rabbit) Specificity

Product No.'s Ig class

1525 IgG1 Specific to the Group B Step carbohydrate. Negative vs. other Strep groups*. Pairs in ELISA ^

II. Purified Preparations

Product No.'s 1525

RabMAB® purified preparations consist of >90% pure rabbit monoclonal antibody which has been purified from culture medium by protein A chromatography or sequential differential precipitations. The final preparation is formulated to a protein concentration of 100 μ g/ml in 0.01 M phosphate buffered saline, pH 7.2 and contains 0.1% sodium azide. Each vial contains 1.0 ml. This product contains no stabilizing proteins and should be stored at 2-8°C until ready for use.

Working dilution must be determined by the user. Suggested starting ranges are 1:10-1:50 for IFA and 1:20-1:200 for ELISA.

THESE PRODUCTS ARE FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES

III. Fluorescein Conjugates

Product No.'s

These MONOTOPETM products consist of purified monoclonal antibody conjugated with high purity isomer I of fluorescein isothiocyanate. Care is taken to ensure complete removal of any free fluorescein from the final product. The final preparation is formulated to an antibody concentration of of 100 μ g/ml in 0.01 M phosphate buffered saline, pH 7.2 containing 0.1% sodium azide plus bovine serum albumin at 10 mg/ml. Each vial contains 1.0 ml. This product should be stored at 2-8°C until ready for use. Avoid repeated freeze-thawing by storing multiple aliquots at -20°C. Applications for these products include direct FA staining of target antigen in a permissive tissue culture system. Working strength must be determined by the user for each specific application but a starting range of 1:5 - 1:20 is recommended. Acetone fixation of the antigen source is recommended prior to staining.

THESE PRODUCTS ARE FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES

Comments: * No cross reactivity with: Streptococci groups A, C, D, F & G + other bacteria.

^ This RabMAB pairs with itself in ELISA to detect the extracted group B carbohydrate antigen.