

MabyRab™ product information for
Mycobacteria LAM

I. Monoclonal Antibody (Rabbit) Specificity

Product No.'s	Ig class	
5179 NA	IgG	Specific for the lipoarabinomannan (LAM) of several Mycobacteria species* Functions in ELISA [^] & IFA.
5180 NA	IgG	Specific for the lipoarabinomannan (LAM) of several Mycobacteria species* Functions in ELISA [^] & IFA.

II. Purified Preparations

Product No.'s	5179
	5180

RabMAB® purified preparations consist of >90% pure rabbit monoclonal antibody which has been purified from culture medium by protein A chromatography or sequential differential precipitations. The final preparation is formulated to a protein concentration of 100 µg/ml in 0.01 M phosphate buffered saline, pH 7.2 and contains 0.1% sodium azide. Each vial contains 1.0 ml. This product contains no stabilizing proteins and should be stored at 2-8°C until ready for use.

Working dilution must be determined by the user. Suggested starting ranges are 1:10-1:50 for IFA and 1:20-1:200 for ELISA.

THESE PRODUCTS ARE FOR RESEARCH USE ONLY NOT FOR USE IN DIAGNOSTIC PROCEDURES

III. Fluorescein Conjugates

Product No.'s	NA
	NA

These MONOTOPE™ products consist of purified monoclonal antibody conjugated with high purity isomer I of fluorescein isothiocyanate. Care is taken to ensure complete removal of any free fluorescein from the final product. The final preparation is formulated to an antibody concentration of 100 µg/ml in 0.01 M phosphate buffered saline, pH 7.2 containing 0.1% sodium azide plus bovine serum albumin at 10 mg/ml. Each vial contains 1.0 ml. This product should be stored at 2-8°C until ready for use. Avoid repeated freeze-thawing by storing multiple aliquots at -20°C. Applications for these products include direct FA staining of target antigen in a permissive tissue culture system. Working strength must be determined by the user for each specific application but a starting range of 1:5 - 1:20 is recommended. Acetone fixation of the antigen source is recommended prior to staining.

THESE PRODUCTS ARE FOR RESEARCH USE ONLY NOT FOR USE IN DIAGNOSTIC PROCEDURES

Comments: * Reactive with: Mycobacterium tuberculosis (6 strains), M. bovis, M. smegmatis, M. leprae, M. avium, M. phlei & M. parafortuitum. See attached table for cross reactivity data.

[^] 5179 as capture + 5180 as detector