

Immunochemicals for Infectious Disease Research www.ViroStat-Inc.com

P.O. Box 8522 Portland, Maine 04104 207-856-6620 207-856-6864 (fax)

MabyRab[™] product information for

Cryptosporidium Oocysts

I. Monoclonal Antibody (Rabbit) Specificity

 Product No.'s
 Ig class

 7641
 IgG1
 Specific to a cyst cell wall glyco-antigen*. Pairs in ELISA to detect cysts ^ also functions in IFA.

 7642
 IgG1
 Specific to a cyst cell wall glyco-antigen*. Pairs in ELISA to detect cysts ^ also functions in IFA.

II. Purified Preparations Product No.'s

RabMAB® purified preparations consist of >90% pure rabbit monoclonal antibody which has been purified from culture medium by protein A chromatography or sequential differential precipitations. The final preparation is formulated to a protein concentration of 100 μ g/ml in 0.01 M phosphate buffered saline, pH 7.2 and contains 0.1% sodium azide. Each vial contains 1.0 ml. This product contains no stabilizing proteins and should be stored at 2-8°C until ready for use.

7641

7642

Working dilution must be determined by the user. Suggested starting ranges are 1:10-1:50 for IFA and 1:20-1:200 for ELISA.

THESE PRODUCTS ARE FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES.

III. Fluorescein Conjugates

Product No.'s

These MONOTOPETM products consist of purified monoclonal antibody conjugated with high purity isomer I of fluorescein isothiocyanate. Care is taken to ensure complete removal of any free fluorescein from the final product. The final preparation is formulated to an antibody concentration of of 100 μ g/ml in 0.01 M phosphate buffered saline, pH 7.2 containing 0.1% sodium azide plus bovine serum albumin at 10 mg/ml. Each vial contains 1.0 ml. This product should be stored at 2-8°C until ready for use. Avoid repeated freeze-thawing by storing multiple aliquots at -20°C. Applications for these products include direct FA staining of target antigen in a permissive tissue culture system. Working strength must be determined by the user for each specific application but a starting range of 1:5 - 1:20 is recommended. Acetone fixation of the antigen source is recommended prior to staining.

THESE PRODUCTS ARE FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES.

Comments: ^ Best ELISA pairs (capture-conjugate): 7641-7642, 7642-7642, 7642-7641 * Periodate sensitive

The antibodies are non-reactive with: Giardia lamblia, Candida albicans & human cells