

**MONOTOPE™** product information for

**Campylobacter jejuni**

**I. Monoclonal Antibody (Mouse) Specificity**

Product No.'s	Ig class	
7711	IgG1	Appears specific to C. jejuni via whole cell ELISA*. Reactive with a ~100 kDa antigen in WB.
7721	IgG1	Appears specific to C. jejuni via whole cell ELISA*. Reactive with a ~100 kDa antigen in WB. Works in LF.
7731	IgG1	Appears specific to C. jejuni via whole cell ELISA*. Reactive with a ~100 kDa antigen in WB.
7741	IgG2a	Appears specific to C. jejuni via whole cell ELISA*. Reactive with a ~100 kDa antigen in WB.

**II. Purified Preparations**

Product No.'s	7711
	7721
	7731
	7741

MONOTOPE™ purified preparations consist of >90% pure mouse monoclonal antibody which has been purified from ascites fluid or culture medium by protein A chromatography or sequential differential precipitations. The final preparation is formulated to a protein concentration of 100 µg/ml in 0.01 M phosphate buffered saline, pH 7.2 and contains 0.1% sodium azide. Each vial contains 1.0 ml. This product contains no stabilizing proteins and should be stored at 2-8°C until ready for use.

Working dilution must be determined by the user. Suggested starting ranges are 1:10-1:50 for IFA and 1:20-1:200 for ELISA.

THESE PRODUCTS ARE FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES

**III. Fluorescein Conjugates**

Product No.'s

These MONOTOPE™ products consist of purified monoclonal antibody conjugated with high purity isomer I of fluorescein isothiocyanate. Care is taken to ensure complete removal of any free fluorescein from the final product. The final preparation is formulated to an antibody concentration of 100 µg/ml in 0.01 M phosphate buffered saline, pH 7.2 containing 0.1% sodium azide plus bovine serum albumin at 10 mg/ml. Each vial contains 1.0 ml. This product should be stored at 2-8°C until ready for use. Avoid repeated freeze-thawing by storing multiple aliquots at -20°C. Applications for these products include direct FA staining of target antigen in a permissive tissue culture system. Working strength must be determined by the user for each specific application but a starting range of 1:5 - 1:20 is recommended. Acetone fixation of the antigen source is recommended prior to staining.

THESE PRODUCTS ARE FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES

**Comments:** \* non-reactive with the following: H. pylori, E. coli (several), Salmonellae (several), Shigellae (several), P. aeruginosa, Yersinia & Citrobacter